<u>THIE UNITHED STATES OF AMIERIOA</u>

TO ALL TO WHOM THESE; PRESENTS; SHALL COME;

Northrup King Co.

CAncreas, there has been presented to the

Secretary of Agriculture

AN APPLICATION REQUESTING A CERTIFICATE OF PROTECTION FOR AN ALLEGED NOVEL VARIETY OF SEXUALLY REPRODUCED PLANT, THE NAME AND DESCRIPTION OF WHICH ARE CONTAINED IN THE APPLICATION AND EXHIBITS, A COPY OF WHICH IS HEREUNTO ANNEXED AND MADE A PART HEREOF, AND THE VARIOUS REQUIREMENTS OF LAW IN SUCH CASES MADE AND PROVIDED HAVE BEEN COMPLIED WITH, AND THE TITLE THERETO IS, FROM THE RECORDS OF THE PLANT VARIETY PROTECTION OFFICE, IN THE APPLICANT(S) INDICATED IN THE SAID COPY, AND WHEREAS, UPON DUE EXAMINATION MADE, THE SAID APPLICANT(S) IS (ARE) ADJUDGED to be entitled to a certificate of plant variety protection under the LAW.

NOW, therefore, this certificate of plant variety protection is to grant UNTO THE SAID APPLICANT(S) AND THE SUCCESSORS, HEIRS OR ASSIGNS OF THE SAID APPLI-CANT(S) FOR THE TERM OF eighteen YEARS FROM THE DATE OF THIS GRANT, SUBJECT TO THE PAYMENT OF THE REQUIRED FEES AND PERIODIC REPLENISHMENT OF VIABLE BASIC SEED OF THE VARIETY IN A PUBLIC REPOSITORY AS PROVIDED BY LAW, THE RIGHT TO EX-CLUDE OTHERS FROM SELLING THE VARIETY, OR OFFERING IT FOR SALE, OR REPRODUCING IT.

IMPORTING IT, OR EXPORTING IT, OR USING IT IN PRODUCING A HYBRID OR DIFFERENT TY THEREFROM, TO THE EXTENT PROVIDED BY THE PLANT VARIETY PROTECTION ACT. United States seed of this variety (1) shall be sold by variety name only as OF CERTIFIED SEED AND (2) SHALL CONFORM TO THE NUMBER OF GENERATIONS THE OWNER OF THE RIGHTS. (84 STAT. 1542, AS AMENDED, 7 U.S.C. 2321 ET SEQ.)

WHEAT

'Coker 9877'

In Testimony Whereof, I have hereunto set my hand and caused the seal of the Plant Tariety Protection Office to be affixed at the City of Washington, D.C. this 30th day of April in the year of our Lord one thousand nine hundred and ninety-three.

City Es Secretary of Agriculture

Attest

Agricultural Marketing Service

Public reporting burden for this collection of information is estimated to average 30 minutes per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Department of Agriculture, Clearance Office, OIRM, Room 404-W, Washington, D.C. 20250; and to the Office of Management and Budget, Paperwork Reduction Project (OMB #0581-0055), Washington, 20250.

U.S. DEPARTMENT OF AGRICULTU AGRICULTURAL MARKETING SERV	JRE /ICE			Application is required in order to
APPLICATION FOR PLANT VARIETY PRO		CERTIFICATE		determine if a plant variety protection certificate is to be issued (7 U.S.C. 2421). Information is held confidential certificate is issued (7 U.S.C. 2426).
NAME OF APPLICANT(S) (as it is to appear on the Certificate)	<u>. </u>	2. TEMPORARY DESIGN		3. VARIETY NAME
Northrup King Co.		CT-84101	19 NOV	Coker 9877
1101 1111 11 11 11 11 11 11 11 11 11 11		CL84010	,	COREL JULI
4. ADDRESS (street and no. or R.F.D. no., city, state, and ZIP)		5. PHONE (Include area		FOR OFFICIAL USE ONLY
P. O. Box 959	. 1			PVPO NUMBER
Minneapolis, MN 55440	İ	(612) 593-73	333	
			- 1	9000222
	1			5 Date
				Ques 9 1990
6. GENUS AND SPECIES NAME 7 FAMI	LY NAME (Botanica			1
Triticum aestivum	Gramineae			G A.M. TP.M.
	or amilioue	•	ŀ	
8. CROP KIND NAME (Common Name)	9. D	ATE OF DETERMINATION		Filing and Examination Fee:
Soft red winter wheat		1988		E 13/3/
TO JETHE ADDITIONAL MANAGERICANT A MECHANISM STORY OF STORY				S Date O. A. O. 1000
10. IF THE APPLICANT NAMED IS NOT A "PERSON," GIVE FORM OF ORGANIZATION ((Corporation, partne	ership, association, etc.)	. 1	R July 9,1990
Corporation				C Certificate Fale:
11. IF INCORPORATED, GIVE STATE OF INCORPORATION	12. DAT	E OF INCORPORATION		‡ <u>\$250.00</u>
Delaware		1976		V Date
				5 Ayou 8, 1993
13. NAME AND ADDRESS OF APPLICANT REPRESENTATIVE(S), IF ANY, TO SERVE IN			RS	
	044 14	orne 2		4
Northrup King Co.		> ^~	t 10%	6 1993
P. O. Box 989 4	TA 5	(252)		
Minneapolis, MN 55440 Washington,	. J-/t-)	PHONE (Inclu	de area code,	_: (612) 593-7305
14. CHECK APPROPRIATE BOX FOR EACH ATTACHMENT SUBMITTED (Follow INSTRU	ICTIONS on reverse)		
a. A Exhibit A, Origin and Breeding History of the Variety.	• *			
b. X Exhibit B, Novelty Statement.				
c. X Exhibit C, Objective Description of Variety.	*			
d. Exhibit D, Additional Description of Variety.				
e X Exhibit E, Statement of the Basis of Applicant's Ownership.				
Seed Sample (2,500 viable untreated seeds). Date Seed Sample in	nailed to Plant Va	riety Protection Office		
g. Kiling and Examination Fee (\$2,150) made payable to "Treasurer of	of the United Stat	es."		
15. DOES THE APPLICANT(S) SPECIFY THAT SEED OF THIS VARIETY BE SOLD BY VAR			SEED? (See	section 83(a) of the Plant Variety
Protection Act.) X YES (If "YES," answer items 16 and 17 below)	_	" skip to item 18 below)		
16. DOES THE APPLICANT(S) SPECIFY THAT THIS VARIETY BE LIMITED AS TO	17. IF "YES" TO	TEM 16, WHICH CLASSES	OF PRODUC	TION BEYOND BREEDER SEED?
NUMBER OF GENERATIONS?	t (₹₽3		991	(er)
X YES NO	E X FOUN	DATION	X REGISTE	RED X CERTIFIED
18. DID THE APPLICANT(S) PREVIOUSLY FILE FOR PROTECTION OF THE VARIETY IN T	HE U.S.?	· · · · · · · · · · · · · · · · · · ·		
	ent Act. Give date:)		
X NO				
19. HAS THE VARIETY BEEN RELEASED, USED, OFFERED FOR SALE, OR MARKETED	IN THE U.S. OR OT	HER COUNTRIES?		
	W THE O.S. ON ON	HER COUNTRIES:		•
YES (II "YES," give names of countries and dates) U.S.A., fall of 1989		•		
□ NO 0.5.A., Tall of 1909				
The state of the s				
The applicant(s) declare(s) that a viable sample of basic seeds of this request in accordance with such regulations as may be applicable.	is variety will b	e furnished with the	application	and will be replenished upon
-				
The undersigned applicant(s) is (are) the owner(s) of this sexually uniform, and stable as required in section 41, and is entitled to prote	reproduced no ection under the	vei plant variety, at	id believe: 42 of the Pl	s) that the variety is distinct, ant Variety Protection Act
Applicant(s) is (are) informed that false representation herein can je				
SIGNATURE OF APPLICANT [Owner(s)]	CAPACITY OR TIT	LE		DATE
Kole VIII Non	Robert	W. Romig		June 22, 1990
1 100. Torres			earch	June 22, 1990
SIGNATURE OF APPLICANT (Owner(s))	CAPACITY OR TIT			DATE

Exhibit A Origin and Breeding History of the Wheat Variety, Coker 9877

Coker 9877 soft red winter wheat is derived from a cross Coker 65-20 4*/Wichita 7-Transfer//Blueboy selection/3/C71-21/Blueboy II made in 1978 at the Coker research facility at Hartsville, South Carolina. C71-21 is a selection from the cross C57-6/Hadden//Blueboy. C57-6, in turn, is a selection from the cross Hardy Red/Taylor. The variety is derived from single plant selection in the F2-F4 generations followed by further selection in the F8 generation.

We grew the F1 through F5 generations at Hartsville, South Carolina where we selected for resistance to <u>Puccinia recondita</u> (leaf rust) and <u>Erysiphe graminis</u> (powdery mildew). In the spring of 1984, we harvested an F5 row (F6 seed) in bulk which we designated C84A77.

On the basis of a performance test in the 1984-85 season at Bay, Arkansas, we advanced this line for further testing in the 1985-86 and 1986-87 season. In 1986-87, we selected individual plants from the experimental variety which was renumbered as CL84101. It has been tested extensively in the Mid-South for the four years (1986 through 1989) and was grown in the USDA Uniform Southern Soft Wheat Nursery for two years (1986-87 and 1987-88). The variety was named Coker 9877 in 1988.

Breeder seed was produced by combining seed from 83 single plant-derived, second-year increases that appeared uniform and stable. This seed source will be maintained for future seed stock purposes. The strip increases combined to produce breeder seed were in the F11 generation and were judged identical in appearance to the original F9 plants selected as representative of the line. The variety does have a low level of variants. These include plants with earlier maturity, taller height, and awned spikes. Frequency of these types has been less than 5/1,000. Coleoptile color is predominantly white/green, but the variety may contain up to 2% purple coleoptiles.

Exhibit B Novelty Statement for the Wheat Variety, Coker 9877

Wheat variety Coker 9877 is most similar to Coker 9733 and Coker 833. It can be differentiated from Coker 9733 on the basis of coleoptile color, maturity, and testweight. Coker 9877 has white coleoptile color vs. purple for Coker 9733. It heads six days later, and has approximately 3.7 pounds per bushel lower testweight than Coker 9733 (see data below).

Coker 9877 can be differentiated from Coker 833 on the basis of resistance to stripe rust, stem rust, and <u>Septoria nodorum</u>. Coker 9877 is susceptible to stripe rust but resistant to stem rust and <u>Septoria nodorum</u>; Coker 833 is resistant to stripe rust but susceptible to stem rust and <u>Septoria nodorum</u>.

Agronomic data for Coker 9877 and Coker 9733 grown for four years in variety trials at Bay, AR.

<u>Variety</u>	<u> 1986</u>	<u> 1987</u>	1988	<u> 1989</u>	<u> Mean</u>						
				• •							
Heading Date (Days in April)											
0-1											
Coker 9877	4/26	4/30	4/28	4/30	4/29						
Coker 9733	4/19	4/24	4/22	4/25	4/23						
$\mathrm{LSD}_{(0.05)}$					1.8 days						
											
Coker 9877	-58	52	57	56	55.8						
Coker 9733	61	59	5 <i>7</i>	59	59.5						
			33		33.3						
LSD _(0.05)				-	2.3 Lb./Bu.						

EXHIBIT C

U.S. DEPARTMENT OF AGRICULTURE AGRICULTURAL MARKETING SERVICE LIVESTOCK AND SEED DIVISION BELTSVILLE, MARYLAND 20706

OBJECTIVE DESCRIPTION OF VARIETY WHEAT (TRITICUM SPP.)

	CON OFFICIAL MET ONLY
NAME OF APPLICANTIS	FOR OFFICIAL USE ONLY
Northrup King Co. ADDRESS (Street and No. or R.F.D. No., City, State, and EIP Code)	9000222
	VARIETY HAME OR TEMPORARY
Attention: R. W. Romig	DESIGNATION
P.O. Box 959	Coker 9877
Minneapolis, MN 55440	
Place the appropriate number that describes the varietal character of this variety in the Place a zero in first box (e-s- 0 8 9 or 0 9) when number is either 99 or less to	
	a y or tess.
I, KIND:	
1 1 = CONNON 2 = OURUM 3 = EMMER 4 = SPELT 5 = POLISH 6 = POL	JLARO 7 = CLUB
2. TYPE,	
ا الشرا ا ≠ soft	3 = OTHER (Specify)
2 1 * SPRING 2 = WINTER 3 = OTHER (Specify) 1 2 = HARD	
2 1 = WHITE 2 = RED 3 = OTHER (Specily)	
3. SEASON - NUMBER OF DAYS FROM EMERGENCE TO:	
, senson - nonsen or only man characters (c)	
2 0 3 FIRST FLOWERING 2 1 0 LAST	Y FLOWERING
(, HATURITY (50% Flowering):	
. MATORITI (30% Flowering):	
HO. OF DAY'S EARLIER THAN	2 = SCOUT 3 = CHRIS
4 = LEMHI	S = NUGAINES 6 = LEEDS 7=Caldwel
0 2 NO. OF DAYS LATER THAN	James To Service To Garages
. PLANT HEIGHT (From soil level to top of head):	
, PCANT NCIONI (From sell level to tap of head):	
0 9 8 cm. High	
U 3 CH, TALLER THAN	e de la companya del companya de la companya de la companya del companya de la co
] = ARTHUR	2 = SCOUT 3 = CHRIS
CM. SHORTER THAN	5 = NUGAINES 6 = LEEDS 7=Caldwel
PLANT COLOR AT BOOTING (See reverse): 7. ANTHER COLORI	
	*** **** **** **** **** **** **** **** ****
1 = YELLOW GREEN 2 = GREEN 3 = BLUE GREEN 1 1 1 = YELLOW	2 = PURPLE
STEM:	
Anthocyanin: 1 = ABSENT 2 = PRESENT Vary bloom: 1:	# ABSENT 2 # PRESENT
Hairiness of last	
internode of rachis: 1 = ABSENT 2 = PRESENT 1 [aternodes: 1 =	
	is almost solid
5 NO. OF HODES (Originating from node above ground) CM. INTER	RHODE LENGTH BETWEEN.FLAG LEAF F BELOW
AURICLES:	
אסאוטנב:	
Anthocyanin: 1 * ABSENT 2 * PRESENT 2 Hairiness: 1 *	ABSENT 2 # PRESENT
	A Company of the Comp
LEAF!	
Flag leaf at 1 = ERECY 2 = RECURVED	OT TWISTED 2 = TWISTED
booting stage: 3 = OTHER (Specify): 1 Flag feat: 1 = N	U
],,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,,	
Hairs of first leaf sheath: 1 = ABSENT 2 = PRESENT Vary bloom of fle	ag leaf sheath: 1 = ABSENT 2 = PRESENT
MM. LEAF WIOTH (First loaf bolow flag load) CH. LFAFE	LENGTH (First leaf below flat leaf):

			· · · · · · · · · · · · · · · · · · ·
3 Density: 1 = LAX	7 = DENSE 3=semi-dense	Shape: 1 = TAPE 2	RING 2 = STRAP 3 = CLAVATE R(Specify)
2 Awnedness: I = Aw	HLESS ? = APICALLY AWHLETED	3 = AWNLETED 4 = AWN	εο
7 Color at maturity: 5	# BROWN 6 # BLACK 7 # OTH	FRED ER (Specity): tan	
0 9 CM. LENGTH		1 2 MM. WIDTH	
12. GLUMES AT MATURI 2 Length: 1 = SHORT 3 = LONG	(CA. 7 mm.) 2 = MEDIUM (CA. 8 mm.)	2 Width: 1 = HARRO 3 = WIDE (7
Shoulder 1 = WANT shape: 4 = SQUA	· · · · · · · · · · · · · · · · · · ·	1 Beak: 1 = OBTUS	(3)
13. COLEOPTILE COLOR	1	14. SEEDLING ANTHOC	YANIN: RECEIVED
1 1 = WHITE 2 = R	ED 3 = PURPLE	1 = ABSENT	2 = PRESENT 8 USDA AMS
15. JUYENILE PLANT GR	ONTH HABIT:	•	☐ JUL 9 - 1990 》 ☐
2 1 = PROSTRATE	2 = SEMI-ERECT 3 = ERE	ccr	Plant Variety L2
16. SEED:			Francoion Ole
Shape: 1 = OVATE 4=Ovate to		1 Check: 1 = ROUNG	DED 2 = ANGULAR
2 Brush: 1 = SHORT	2 = MEDIUM 3 = LONG	Brush: 1 = NOT C	COLLARED 2 = COLLARED
Phenol reaction (See instructions):	1 = IVORY 2 = FAWN 3 = LT. BROW 4 = BROWN 5 = BLACK	WH .	•
5 Color: 1 = WHITE	2 = AMBER 3 = RED 4 = PURPLE	5 = OTHER (Specily)	medium brown
0 7 MM. LENGTH *	. 0 3 мм. wютн	2 8 GM. PER 1000	*variable due to tendency to set 3 to 5 seed/spikelet
17. SEED CREASE:			•
2 Vidth: = 60% OR L	ESS OF KERNEL 'WINOKA'	1 71	R LESS OF KERNEL 'SCOUT'
2 = 80% OR LI	narrow to mid wide	2 = 35% 0	R LESS OF KERNEL 'CHRIS' mid-deep
3 = HEARLY	VS MIDE YS KENNEF , FEMHI.	3 = 50 % 01	R LESS OF KERNEL 'LEMHI'
18. DISEASE: (0 = Not Test	ed, 1 = Susceptible, 2 = Resistant) 3	dependent	
2 STEM RUST (Races)	2 LEAF RUST (Races)	race dependent STRIPE RUST (Recee)	O LOOSE SHUT
3 POWDERY MILDEW Susceptible SE rad	ce; Resistant to Mid-South	Tace OTHER (Specify) S	eptoria leaf & glume blotch
	d, 1 = Susceptible, 2 = Resisions)		
O SAWFLY.	O APHID (Byon)	O GREEN BUG	O CEREAL LEAF BEETLE
OTHER (Specily)	HESSIAN FLY	GP - A	В С
	RACES:	O 1 E	
20. INDICATE WHICH VARIE	TY HOST CLOSELY RESEMBLES THAT S	UBMITTED:	
CHARACTER	HAME OF VARIETY	CHARACTER	HAME OF VARIETY
Plant tillering	Coker 833	Seed size	Coker 833
Leaf size	Coker 833	Seed shope i	Coker 833
Leaf color	Coker 833	Coleoptile elongation	Coker 833
Leaf corriage	Coker 833	Seedling pigmentation	Coker 833
·			

INSTRUCTIONS

GENERAL: The following publications may be used as a reference aid for the standardization of terms and procedures for completing this form:

- (a) L.W. Briggle and L. P. Reitz. 1963. Classification of Triticum Species and Theat Varieties Grown in the United States. Technical Bulletin 1278, United States Department of Agriculture.
- (b) W.E. Valls, 1965, A Standardized Phenol Method for Testing Wheat Seeds for Varietal Purity, contribution No. 28 to the handbook of seed testing prepared by the Association of Official Seed Analysts. (See attachment.)

Milling and Baking Characteristics of Coker 9877*

Advanced Nursery Evaluations Average of 1986-88

Quality Parameter	Coker 9877	Coker 916	Coker 9323
Milling Quality Score	91.3	93.9	103.9
Baking Quality Score	87.8	84.3	94.7
Test weight (kg/h1)	77.9	78.0	78.0
Softness Equivalent	53.0	55.3	59.9
Flour Yield	70.5	70.2	72.7
Ash	.39	.36	.31
Flour Protein	10.0	10.4	10.8
Micro AWRC	53.5	54.1	52.0
Cookie Diameter	17.4	17.2	17.7
Top Grain	3.7	3.0	3.3

^{*}Milling and baking quality tests conducted by USDA, ARS Soft Wheat Quality Lab; Wooster, Ohio.

Also in 1986-87 and 1987-88 USDA coordinated Uniform Southern Soft Wheat Nurseries, Coker 9877 ranked 16th/34 and 19th/32, respectively.

DATA RANKED ACCORDING TO COMBINED QUALITY SCORE

1987 CROP

UNIFORM SOUTHERN NURSERY STANDARD = 340, TYLER

WHEAT AND MILLING DATA

LAB NO.		ENTRY	MILLING QUALITY SCORE	BAKING QUALITY SCORE	COMBINED QUALITY SCORE	TEST VT.	BREAK FLOUR YIELD	ST.GR. FLOUR YIELD	RED PASSES	FRIABILITY	Ε.\$.Ι.	MILLABILITY
			*****	*******	***************************************							.1
347		FL 7924-D13-H14	111.A	110.7A	110.A	59.9	33.70	11.8	1	30.5	8.1	131.8
339	FR	PIONEER 2555 (XY 546)	108.4	115.5A	108.A	59.2	42.1	77.3	1	30.4	9	122.3
338	8EX	STEELE (NA-SV-76-59)	105 A	113.5Å	105 A	58.4Q	37.4	77.7	1	30.8	8.5	120.3
369	860	NA-SV-84-160	104.A	109.5A	104.A	61.1	37.5	11.5	1	29.5	9.7	117.1
337	AQ	FLORIDA 302	107.A	102.94	102.A	60.9	33.4Q	77.3	7	29.8	8.6	123.5
***		. BENCHWARK	100.A	112.14	100.A	61.6	35.6*	75.3	7	28 Q	10.4	109.9
348	20	FL 7927-621	102.A	100.14	100.A	81.3	35 Q	78.9	7	28.7	10.2	113.4
340	8,J	TYLER	100 A	100 A	100 A	60.1	37.3	76.3	7	29	10.4	109.9
111		STANDARD	100 A	100 A	100 A	60.1	37.3	76.3	7	29	10.4	109.9
361	BGI	GA 781197	100 A	103.64	100 A	59.7	38.2	76.2	7	29.3	9.9	109.8
344	BFK	AR 116-6	99 B	102.1A	99 B	51.3	33.20	76.4	7	28.9	10.1	109
367	BGK	ND 73025-51	98.78	111.43	98.7B	58.20	38	11.3	1	29.2	10.1	109.4
341	BFR	COKER 9733 (COKER 84-33)	97.28	99.98	97.2B	81.7	30.80	15.2*	7	29	10.8	108.8
362	BGD	TX-79-30	97.18	110 A	97.1B	81.5	38.8	75.5	7	27.7 Q	11.5#	102.8
345	8FL	AR 322-21-3	101.A	96.9 8	96.98	60.5	31.10	77.2	7	29.3	9.9	115
354	850	COKER 86-29	95.38	95.3 8	95.38	62.1	33.8Q	75.6	1	28.5 *	10.9	193.1
352	BFS	COKER 86-17	95.28	103.34	95.28	60.9	37.9	75.8	1	28.9	12 1	100.3*
350	BFQ	CL 840101 - Coker 9877	95.18	113 A	95.18	80.7	31.8Q	78.1	7	28.5	11.3	103.8
353	8FT	COKER 86-26	95.1B	96.2 8	95.18	82.2	31.7Q	75.8	1	28.4 *	10.8	102.1
370	BGP	NA-SV-84-313	94.4C	94.7 C	94.4C	58.8Q	31.8Q	76	1	28.7	10.4	104.8
384	8EC	TX-78-7303	99.58	91.7 C	91.7C	59.4	37.3	78.1	8	28.4 *	11.6	109.8
343	BGN	HA-SY-17-42-25	91.5C	94.4 €	91.5C	80.8	37.4	75.1	8	26.8 Q	12.4€	94.3 *
365	BGN	ND 75191-80	103.A	90.4 C	90.40	59.1*	34.30	17.2	1	29.4	9.4	118.3
363	8ED	TX-79-19-1	89 D	89.3 D	89 D	59.2	35.9	75.3*	8	26.9 Q	12.34	92.6 *
346	BGA	SC 810779	88.SD	90.5 C	88.50	62	31.90	74.9*	8	25.9 Q	12.80	90.8 Q
366	86J	ND 73019-28	87.70	94.7 C	87.7D	59.7	37.5	74.9*	8	26.6 Q	12.99	88.9 Q
360	Beh	GA 781178	90 C	86.4 D	86.40	59.3#	36.8	74.81	1	27.5 Q	12.50	93.7 *
368	BGL	ND 73065-03	85.7D	110.94	85.70	60.2	39	75.5	8	26.5 Q	12.14	84.1 Q
342	BGE	VA 82-52-64	86.90	84.1 E	84.1E	60	39.4	75 *	7	27.5 Q	11.6#	86.2 Q
349	BGF	FL 72185A101-5-9	102.A	83.2 E	83.2E	62.2	32.7Q	76.4	. 1	28.9	10.1	114.2
355	8F¥	COKER 88-31	101.A	83 E	83 E	81.2	12.90	76.7	7	29.3	9.5	114.3
351	BFN	AL 840169	97.5B	82.5 E	82.58		36.11	76.4	7	28.4 *	10.8	108.1
	BFN	AT 83WHR6208	97 8	78.8 F	78.8F	61.1	34.8Q	76	8	27.9 Q	11.4*	105.1
357	868	SC 820556	91.5C	11.2 F	77.2F	61.5	31.BQ	75.9	7		11.3	98.8 *
358	86C	SC 820885.	100.4	48 F	46 F	61.3	21.90	11.6	1		3.7	117.9
359	866	GA 781159	88.6D	25.7 F	25.7F	63.2	19.8Q	-78.1	7	26.5 Q	11.5#	92.7 *

DATA RANKED ACCORDING TO CONBINED QUALITY SCORE

1.987 CROP

UNIFORM SOUTHERN NURSERY STANDARD = 340, TYLER

STRAIGHT-GRADE FLOUR

:					
LAB		FLOUR			
y NO.		HEA		DIAMETER	GRAIN
` ^·	\$	\$	\$	CM.	
347		.35	48.7	18.2	δ
339		.38	50.8	18.45	5
338		.41	50.1	18.43	\$
369	8.03	,39	53 #	18.22	5
337	8.76	.37	48.5	17.94	i i
111	8.899#	.35	51.3	18.35	7
348	9.31 Q	.38		17.95	6
340	8.38	.39	50.7	17.91	5
	8.36 1	.39	50.7	17.91	5
361	9.42 Q	.4	50.9	18.09	5
344	9.91 Q	.4	49.5	18.03	7
367	8.66	.42 *		18.18	8
341	9.7 Q	.38	17.1	17.94	5
362	9.22 *	.38	51.9	18.33	5
345	9.04 *	.39	49.4	17.83	5
354	10.060		49.4	17.87	3
352	9.65 Q	41	49.8	18.05	2
350	9.02 *	.4	50.6	18.29	8
353	10.660	.41	19.8	17.95	1
370	9.07 *	.41	50.3	17.82	5
364	9.41 Q	.36	51.8	17.84	3
343	8.95 *	.39	51.8	17.86	7
365	8.15	.38	50.9	17.67	5
363	9.27 *	-41	52.2	17.79	3
348	10.45Q	.39	51.5	17.88	5
385	8.37	.41	52.2	17.85	3
360		. 4	52.7 \$	17.78	5
368	9,11 *	.46 Q	51.6	18.36	5
342	8.84	.46 Q	53.8 *	17.7	3
349	9.95 Q	.37	50.3	17.81*	4
355	9.18 *	.39	49.8	17.52*	5
65 351	10.27Q	.39 .38	50.6	17.63*	3
355 351 356 357	9.75 Q		91: <i>1</i> ,	17.55*	2 .
. 441	10.204	. 43 *		17.59*	5
358	9.82 Q	.39	•	15.92Q	Į.
359	9.9. Q	.43 *	59.1 Q	18.59Q	2
	٠,				

DATA RANKED ACCORDING TO CONSINED QUALITY SCORE

1988 CROP

UNIFORM SOUTHERN NURSERY STANDARD = 88337, FLA 302

WHEAT AND WILLING DATA

LAB No.		ENTRY	HILLING QUALITY SCORE	BÀKING QUALITY SCORE	COMBINED QUALITY SCORE	TEST YT:	BREAK FLOUR YIELD	ST.GR. FLOUR YIELD	RED PASSES	FRIABILITY	E.\$.I.	MILLABILITY
***		BENCHWARK	102.Å	100.7A	100.A	61.6	35.8	76.3*	1	.29 Q	10.40	123.7
***		STANDARD	100 A	100 A	100 A	61.3	30.3	11.9	1	29.9	8.6	123.7
337	2	FLA 302	100 A	100 A	100 A	61.3	30.3	11.8	1	29.3	8.8	123.7
342		PIONEER 2555	98.88	114.2A	98.88	81.8	34.5	11.1	7	30	8.8	117.9
347	12	NASV 84-160	97.18	97.2 8	97.18	62.5	31.3	18.2	1	29.4 *	9	115.9
352		ND 73025-51	98.48	103.6A	96.48	58.90	31	78.5	7	29.7	8.8	119.1
363	28 .	COKER 87-13	98 8	95.5 B	95.58	82.8	32,5	17.2	1	29.3 #	9.3	118.8
355	20	FL 7927-G26	98.68	95.4.B	95.48	61	31.1	78.3	7	30	8.5	120.8
338	3	FLA 303	95.18	105 A	95.18	δ2	30.1	17.4	7	28.8 2	9.5 \$	113.5
349	14	TX 79-30	94.5C	106.7A	94.50	82.7	33.5	76.3	1	28.9 Q	10.7Q	108.8≇
385	30	VA 85-52-24	94.10	97 8	94.1C	61.4	26.70	11.4	7	29.5 #	9.8 *	114.3
364	29	GA 801310	93.4C	96 8	93.4C	80.8*	27.5Q	18.1	1	28.3 Q	10.1	113.7
386	31	VA 85-52-34	93.2C	100.7Å	93.2C	61.5	31.3	76.9	7	28.4 Q	10.1*	109.4*
360	25	CL 850655	93 C	93 C	93 C	52.4	30.7	16.5#	1	28.5 Q	10.7Q	108.5*
346	11	GA 781197	92.60	109.1A	92.8C	61.4	33.8	78.8	1 -	29 Q	3.4 \$	108.5*
338	1	TYLER	92.5C	99.88	92.5C	80.5#	29.8	75.4*	7	28.6 Q	10.6Q	110.4*
358	21	FL 1927-G29	100.1	92.4 C	92.4C	82.8	28.8	78.4	1	29.8	8.5	124.1
359	24	GA 80078-P1988	96.5B	91.9 C	91.90	61.9	32.8	76.48	7	29.2 *	10.4ହ	114.2
362	21	COKER 87-12	95.58	89.2 D	89.20	82	31.7	17.5	7	28.9 Q	9.3	113.1*
351	18	ND 75191-80	97.18	88.8 D	88.8D	59.40	30.9	11.5	1	29,2 *	9	119.9
341	8	COKER P9877	88.70	90.1 C	88.70	81.8	27.4Q	78.1	7	28.2 ♀	11.3Q	103.43
350	15	TX 79-19-1	90.90	88.6 D	88,6D	81.5	30.9	76.3*	8	27.7 Q	10.99	105.2*
340	5	BAYLES	90.70	88 0	88 0	81.1	28.9*	78.8	7	27.5 Q	10.7Q	106.9*
353	18	ND 73085-03	88.2D	89.3 D	86.20	80.1Q	31.3	75.9Q	1	27.4 Q	10.4Q	97.4 Q
348	13	NASW 84-313	86.70	84.9 E	84.98	50.9	26,4Q	16.42	1	28.3 Q	11,29	101.19
339	¥.	NASW77-42-25	90.10	84.8 E	84.88	81.2	31	75.80	7	27.7 Q	11.48	103.9*
367	32	KY 83-27	84 E	107.1A	84 E	60.2*	31.8	15 Q	7	•	11.90	92.9 ફ
354	19	AR 28415	82.6E	79.5 F	79.5F	62.7	29.7	74 Q	7	26.7 Q	12.69	88.5 ପ୍
357	22	FLORIDA 301H6	93.9C	17.6 F	17.68	62.5	28 Q	77.1	7	,	10 *	112.1*
343	8	SC 820558	85.20	74.2 F	74.2F	62.8	28.1Q	76.3*	1	27.4 Q	11.3Q	94.1 Q
	26	C 86-27	87.80	72.4 F	72.4F	63.7	21.60	78.4	7		11,10	99.1 Q
345	10	COKER 86-31		. 10.9 F	70.9F	62.8	28.6Q	76.8	1	•	9.5 *	104.4*
	23	FL 85T¥152-1	88.80	63.9 F	63.9F	63	22.2Q	11	1	•	9.9 *	105.9*
344	9	SC 820886	87.60	42.4 F	42.4F	62.8	18.90	11	1	27.1 Q	9.4 \$	106.3*

DATA RANKED ACCORDING TO COMBINED QUALITY SCORE

1988 CROP

UNIFORM SOUTHERN NURSERY STANDARD = 88337, FLA 302

STRAIGHT-GRADE FLOUR

LAB	FLOUR	FLOUR	MICRO	COOKIE	TOP
Ю.	PROTEIN	ASH	A.W.R.C.		GRAIN
	\$	1	\$	ÇW.	
***	8.899*	.35	51.3	18.35	1
***	8.23	.38	51.8	18.3	5
337	8.23	.38	51.8	18.3	5
342	8.46	,41 *	51.8	19.07	8
347	8.3	.42 Q	52.8	18.28	8
352	8.65	.42 Q	49.9	18.35	\$
363	8.65	.39	52.2	18.24	4
3,55	9.27 Q	.41 *	51.1	18.24	4
338	9 🗜	.4 1	52.3	18.51	6
349	9.38 Q	.39	51.8	18.56	5
365	8.84 \$.4 4	49.6	18.2	4
364	9.38 Q	.37	51.5	18.28	5
388	8.36	.4 *	50.3	18.26	8
360	8.76 *	.39	52.9	18.22	5
346	8.89 \$.43 Q	51.8	18.58	5
336	8.09	.38	51.5	18.27	4
358	8.98 \$.39	51.1	18.14	5
359	8.52	.37	53.3	18.19	7
382	8.76 *	.41 *	51.7	18.07	4
351	7.11	.38	52.5	18.01	6
341	8.33	.41 *	51.3	18.04#	5
350	8.8 \$.39	53.3	18.13	i.
340	9.69 Q	.39	52	18.13	¥ .
353	8.94 *	.43 Q	51.3	18.07	5
348	9.42 Q	.42 Q	52.4	18.05*	5
339	8.45	.38	53.1	18 ‡	5
367	8.42	.42 Q	51.3	18.47	8
354	9.49 Q	.4	52.8	17.94*	3
357	9.87 Q	.39	51.2	17.85#	4
343	11.1 Q	.44 Q	53.5 \$	17.97*	5
361	8,95 *	,43 Q	53.5 *	17.75Q	į.
345	8.95 *	.43 Q	53.2	17.7 Q	5
358	10 Q	.41	54.7 *	17.680.	Ĭ
344	9.16 Q	.42 Q	59.2 Q	17.280	.5
J44	3.10 V	145.4	9316. V	arrest .	•

Exhibit E Statement of the Basis of Applicant's Ownership

Wheat variety Coker 9877 was developed by the Northrup King Cocereals breeding staff from germplasm sources cited in Exhibit A of this application. Northrup King Co. believes that the variety is novel as defined in the Plant Variety Protection Act and, therefore, that Northrup King Co. is the sole owner of the variety.